arXiv:1207.2150v1 [cond-mat.str-el] 9 Jul 2012

Mott transitions in the Hubbard model with spatially-modulated interactions

Akihisa Koga and Takamitsu Saitou
Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan

Atsushi Yamamoto
RIKEN, Advanced Institute for Computational Science, 7-1-26,
Minatojima-minami-machi, Chuo-ku, Kobe, Hyogo 650-0047, Japan

(Dated: July 11, 2012)

We study two-component fermions in optical lattices with spatially alternating on-site interactions by means of dynamical mean-field theory. Calculating a quasiparticle weight, double occupancy, and order parameters for each sublattice, we discuss low-temperature properties in the system. When the repulsive interactions alternate, the magnetically ordered state is realized at half filling, while the superfluid state is realized in the attractive case. Furthermore, we find that the metal-insulator transition occurs against magnetic and superfluid fluctuations when the repulsive and attractive interactions alternate. In the doped system, the commensurability emerges due to the repulsive interactions and the insulating state is realized down to the quarter-filling.

I. INTRODUCTION

Recent extensive experimental and theoretical investigations on ultracold gases have providing a variety of interesting topics. Typical examples are ultracold atoms loaded on optical lattices with tunable and controllable parameters,[1] where many remarkable phenomena have been observed such as the Mott transitions[2–4] and BCS-BEC crossover.[5–7] Recently, the spatial modulation of the atom-atom contact interaction has been realized in ^{174}Yb gas system,[8] which stimulates further theoretical investigations on particle correlations in the ultracold atomic systems.

One of the interesting questions is how the spatial modulation in the interactions affects low-temperature properties in fermionic optical lattice systems. Here, we consider the optical lattice with alternating on-site interactions as a simplest model. This should capture the essence of the systems with the modulated interactions with a finite period. In our previous study,[9] we have clarified that in the half-filled system, the Mott transitions occur when the magnitudes of two interactions increase. However, it is naively expected that magnetically ordered state or superfluid state is stabilized in the system only with repulsive or attractive interactions. Therefore, it is necessary to discuss whether or not the Mott transition is indeed realized against magnetic and superfluid fluctuations. It is also interesting to clarify how the alternating interactions affects low-temperature properties in the doped system. This should be important in discussing low-temperature properties in realistic optical lattice systems where the local particle density varies due to the existence of the harmonic potential.[3,4,10,11] Moreover, the system with alternating interactions can be regarded as the natural extension of the two-band model,[12–14] which has been studied in condensed matter physics. Therefore, it is instructive to systematically study how low-temperature properties are affected by the spatial modulation of the on-site interactions.

In this paper, we consider the Hubbard model with alternating on-site interactions to discuss how particle correlations affect low-temperature properties. To this end, we make use dynamical mean-field theory (DMFT)[15–19] with a continuous-time quantum Monte Carlo (CTQMC) method.[20] This combined method has an advantage in dealing with magnetically ordered and superfluid states on an equal footing in the strong coupling region. Thus, we will show how such states are affected by alternating interactions. Furthermore, we discuss the stability of the metal-insulator transition in the system.

Our paper is organized as follows. In Sec. II, we introduce the model Hamiltonian and briefly summarize our theoretical approach. In Sec. III, we demonstrate how the magnetically ordered and superfluid states are realized at half filling. It is clarified that the metal-insulator transition occurs in the system with both repulsive and attractive interactions. We also discuss the effect of the hole doping to clarify the role of the repulsive interactions in Sec. IV. A brief summary is given in the last section.

II. MODEL AND METHOD

We consider two-component fermions in an optical lattice with alternating on-site interactions, which should be described by the following Hubbard Hamiltonian on a bipartite lattice,

$$\begin{aligned} H &= -t\sum_{\langle ij\rangle\sigma} c_{i\sigma}^\dagger c_{j\sigma} - \sum_{i\sigma}\left(\mu+h\sigma\right)n_{i\sigma} \\ &+ \sum_{i\in A} U_A\left(n_{i\uparrow}-\frac{1}{2}\right)\left(n_{i\downarrow}-\frac{1}{2}\right) \\ &+ \sum_{j\in B} U_B\left(n_{j\uparrow}-\frac{1}{2}\right)\left(n_{j\downarrow}-\frac{1}{2}\right), \end{aligned} \tag{1}$$

where $c_{i\sigma}^\dagger (c_{i\sigma})$ creates (annihilates) a fermion at the ith site with spin σ and $n_{i\sigma} = c_{i\sigma}^\dagger c_{i\sigma}$. t is the hopping integral and U_α is the on-site interaction at the sublattice $\alpha(= A, B)$. μ is the chemical potential and h is the magnetic field. We note that in our model Hamiltonian eq. (1), $\langle n\rangle = 1$ (half filling) is realized when $\mu = h = 0$, where $n = \sum_{i\sigma} n_{i\sigma}/N$ and N is the total number of sites. In the case, there exists the symmetry, in addition to the trivial symmetry axis ($U_A = U_B$). By applying the particle-hole transformation[21] $c_{i\uparrow} \to \tilde{c}_{i\uparrow}$ and $c_{i\downarrow} \to (-1)^i \tilde{c}^\dagger_{i\downarrow}$ to the model Hamiltonian with (U_A, U_B, μ, h), we obtain the model Hamiltonian with $(-U_A, -U_B, h, \mu)$. Therefore, in the case with $\mu = h = 0$, the Hubbard models with (U_A, U_B), (U_B, U_A), $(-U_A, -U_B)$ and $(-U_B, -U_A)$ are identical. In the repulsive model ($U_A > 0$ and $U_B > 0$), a magnetically ordered state expected naively is equivalent to the superfluid state in the attractive model ($U_A < 0$ and $U_B < 0$). We also note that the particle number imbalance ($n_A \neq n_B$), which may be regarded as the density wave state, generally appears since the system has a two-sublattice structure, where $n_\alpha = 2\sum_{i\in\alpha,\sigma} n_{i\sigma}/N$.

The Hubbard model with alternating interactions has been studied in one-dimensional systems, where the possibility of anomalous metallic states is discussed.[22] In the paper, we deal with the infinite-dimensional Hubbard model to discuss how alternating interactions affect low-temperature properties. To clarify this issue, we employ the DMFT method.[15–19] In DMFT, the lattice model is mapped to an effective impurity model, where local particle correlations are taken into account precisely. The lattice Green's function is then obtained via self-consistent conditions imposed on the impurity problem. The treatment is formally exact in infinite dimensions, and even in three dimensions, DMFT has successfully explained interesting physics such as the Mott metal-insulator transition.

When DMFT is applied to the system with a sublattice structure, the Green's function is given as,[23]

$$\hat{G}(\mathbf{k}, z)^{-1} = \hat{G}_0(\mathbf{k}, z)^{-1} - \hat{\Sigma}(z), \tag{2}$$

with

$$\hat{G}_0(\mathbf{k}, z) = \begin{pmatrix} (z+h)\hat{\sigma}_0 + \mu_A\hat{\sigma}_z & -\varepsilon_{\mathbf{k}}\hat{\sigma}_z \\ -\varepsilon_{\mathbf{k}}\hat{\sigma}_z & (z+h)\hat{\sigma}_0 + \mu_B\hat{\sigma}_z \end{pmatrix}^{-1}, \tag{3}$$

and

$$\hat{\Sigma}(z) = \begin{pmatrix} \hat{\Sigma}_A(z) & 0 \\ 0 & \hat{\Sigma}_B(z) \end{pmatrix}, \tag{4}$$

where $\hat{\sigma}_0$ is the identity matrix, $\hat{\sigma}_z$ is the z-component of the Pauli matrix, $\mu_\alpha = \mu - U_\alpha/2$ and $\epsilon_{\mathbf{k}}$ is the dispersion relation for the bare band. $\hat{\Sigma}_\alpha(z)$ is the self-energy for the αth sublattice in the Nambu formalism. The local lattice Green's function is obtained as, $\hat{G}_\alpha(z) = \int dk \hat{G}_{\alpha\alpha}(k, z)$. Here, we use a semicircular density of states, $\rho(x) = \frac{2}{\pi D}\sqrt{1 - \left(\frac{x}{D}\right)^2}$ where D is the half-bandwidth, which corresponds to an infinite coordination Bethe lattice. The self-consistency equation is then given by

$$\hat{\mathcal{G}}_\alpha(z) = z\hat{\sigma}_0 + \mu_\alpha\hat{\sigma}_z - \left(\frac{D}{2}\right)^2 \hat{\sigma}_z \hat{G}_{\bar{\alpha}}(z)\hat{\sigma}_z, \tag{5}$$

where $\hat{\mathcal{G}}_\alpha(z)$ is the non-interacting Green's function of the effective Anderson impurity model for the αth sublattice.

There are various numerical methods to solve the effective impurity problem such as the iterative perturbation theory[24,25] and the exact diagonalization.[13] Here, to discuss low-temperature properties quantitatively, we make use of the CTQMC method.[20] This technique has recently been developed[20] and has successfully been applied to general classes of models such as Hubbard model,[26–28] periodic Anderson model,[29,30] Kondo lattice model[31] and Holstein-Hubbard model.[32] Here, we use the hybridization-expansion version of the CTQMC method[26] extended to the Nambu formalism.[33] This allows us to directly access the superfluid state at low temperatures. In our CTQMC simulations, we measure normal and anomalous Green's functions on a grid of one thousand points. Furthermore, we also use the numerical renormalization group (NRG) method[34] to discuss the stability of the normal metallic state complementary. In the NRG, one discretizes the effective bath on a logarithmic mesh by introducing a discretization parameter Λ. The resulting discrete system can be mapped to a semi-infinite chain with exponentially decreasing couplings, which allows us to access and discuss properties involving exponentially small energy scales.[35] To ensure that sum rules for dynamical quantities are fulfilled, we use the complete-basis-set algorithm.[36–38] We observe that obeying the sum rules is mandatory to properly describe low-energy properties of the system away from half filling. In the NRG calculations, we use a discretization parameter $\Lambda = 2$ and keep four thousands states at each step.

In this paper, we use the half-bandwidth D as the unit of energy. To discuss low-energy properties of the system with alternating interactions systematically, we calculate quasiparticle weight $z_\alpha = (1 - \partial\mathrm{Re}\Sigma_\alpha(\omega)/\partial\omega)^{-1}$, double occupancy $D_\alpha = \langle n_{\alpha\uparrow} n_{\alpha\downarrow}\rangle$, magnetization $m_\alpha = (n_{\alpha\uparrow} - n_{\alpha\downarrow})/2$, and pair potential $\Delta_\alpha = \langle c_{\alpha\uparrow} c_{\alpha\downarrow}\rangle$ for each sublattice. When the CTQMC method is used as an impurity solver, we calculate the quantity $z_\alpha = (1 - \mathrm{Im}\Sigma_\alpha(i\omega_0)/\omega_0)^{-1}$ as a quasiparticle weight at finite temperatures, where $\omega_0 = \pi/\beta$. Furthermore, by applying the maximum entropy method[39–41] to the Green's functions, we deduce the spectral functions, which allows us to discuss static and dynamical properties of the systems.

III. RESULTS AT HALF FILLING

We first discuss low-temperature properties in the Hubbard model at half filling ($n = 1$).[9] When the signs of two interactions are the same ($U_AU_B > 0$), the ordered ground-state is naively expected at zero temperature. Namely, the magnetically ordered state is stabilized in the repulsive case, while the superfluid state is in the attractive case. Here, we consider the Hubbard model with repulsive interactions since the attractive model at half-filling is equivalent under the particle-hole transformation, as discussed above. To clarify how the magnetically ordered state is affected by alternating on-site interactions at low temperatures, we calculate the staggered magnetization $m_{AF} = \sum_i (-1)^i m_i/N$. The obtained results at $T/D = 0.05$ are shown in Fig. 1. In the weak coupling region, the paramagnetic metallic state is realized with $m_{AF} = 0$. Increasing the interactions at both sublattices, the staggered magnetization is induced and the phase transition occurs to the magnetically ordered state. In the state, the magnitude of magnetization is almost uniform in the system even with alternating interactions ($U_A \neq U_B$) although the Green's functions strongly depend on the sublattices. Further increase in the interactions leads to another phase transition to the paramagnetic state since thermal fluctuations are relatively enhanced in comparison with magnetic fluctuations. Since the effective intersite coupling should be scaled as $t^2/(U_A+U_B)$, the phase boundary is almost linear ($U_A + U_B = const.$) at finite temperatures, as shown in Fig. 1. It is expected that the magnetically ordered region is extended at lower temperatures.



FIG. 1: (Color online) Density plot of the spontaneous magnetization in the system with $T/D = 0.05$.

On the other hand, when the signs of interactions are different from each other ($U_AU_B < 0$), ground state properties should be nontrivial. In this case, a nonmagnetic ground state is expected since the cooperative phenomenon is hard to be realized in the framework of DMFT. In fact, we could not find any signature of the ordered state down to $T/D = 0.02$. To discuss the stability of the normal metallic state, we calculate the quasiparticle weights for both sublattices with a fixed ratio $U_B/U_A = -2$ at $T/D = 0.02$, as shown in Fig. 2. When the interactions are turned on, the quasiparticle weights z_A and z_B decrease from unity in slightly different way reflecting the difference of on-site interactions. Namely, a strong renormalization appears in the sublattice B. Further increase in the interactions induces the jump singularity in both curves around $U_A/D \sim 1.7$. Then the double occupancy approaches zero (half) at the sublattice $A(B)$, as shown in the inset of Fig. 2. These imply that the first-order phase transition occurs to the insulating state, where the singly occupied states with spin are realized at the sublattice A, while the empty or doubly occupied states are equally realized at the other. Therefore, we conclude that the Mott and pairing transitions simultaneously occur in the sublattices A and B, respectively.



FIG. 2: (Color online) Quasiparticle weight for each sublattice as a function of U_A/D with a fixed ratio $U_B/U_A = -2$. Solid lines with circles are obtained from DMFT with the CTQMC method at $T/D = 0.02$. Dashed lines with squares are obtained from DMFT with the NRG method. Inset shows the double occupancy in the system.

The density of states at each sublattice at $T/D = 0.02$ is shown in Fig. 3. It is found that two kinds of broad peaks appear in the high energy region. One of them is located around $\omega \sim \pm U_\alpha/2$, which forms the Hubbard gap due to the local Coulomb interactions. The other broad peak around $\omega \sim \pm U_{\bar{\alpha}}/2$ is induced by the on-site interaction at nearest neighbor sites although it may be invisible on this scale for the density of states ρ_A. When $U_A/D = 1.75$, the sharp quasiparticle peaks for both sublattices appear in the vicinity of the Fermi level, which implies that the system is in the metallic state close to the Mott transition point. On the other hand, when $U_A/D = 2.0$, the gap structure appears in both sublattices, which means that the insulating state is induced by the Mott and pairing transitions. Therefore, we conclude that the single metal-insulator transition indeed occurs in the system with alternating interactions ($U_AU_B < 0$).

To examine the nature of the phase transition at zero temperature, we also use the NRG method as an impu-



FIG. 3: Solid (dashed) lines represent the density of states for the sublattice $A(B)$. Upper (lower) panel shows the results for the metallic state with $U_A/D = 1.75$ (the insulating state with $U_A/D = 2.0$) in the system with $U_B/U_A = -2$ at $T/D = 0.02$.

rity solver. In the weak coupling region, the NRG results are in good agreement with the CTQMC results, as shown in Fig. 2. However, increasing the interactions, the quasiparticle weights do not have the jump singularity, and smoothly reach zero at the same critical point ($U_A/D \sim 2.1$). This suggests the existence of the second-order quantum phase transition in the half-filled Hubbard model with alternating interactions, which is similar to the nature of the conventional Mott transition in the paramagnetic state.

By performing similar calculations, we obtain the zero temperature phase diagram, as shown in Fig. 4. When U_A and U_B are positive (negative), the magnetically ordered (superfluid) ground state should be realized. In this case, the Mott (pairing) transition never occurs. By contrast, when the signs of the interactions are different ($U_AU_B < 0$), any ordered states should not be stabilized and the metal-insulator transition occurs. We find that the obtained critical points are in good agreement with the phase boundary $U_AU_B = -(3D)^2$ obtained from the linearized DMFT method.[42] The Mott insulating state is stabilized in the strong coupling region, where the singly occupied states are realized in the sublattice with the repulsive interaction, and the empty or doubly occupied states in the other. In the following, we discuss how alternating interactions affects low-temperature properties in the doped system.



FIG. 4: Ground-state phase diagram of the half-filled Hubbard model with alternating interactions. Solid lines represent the phase boundaries obtained from the linearized DMFT method[42] and circles are obtained from DMFT with the NRG method.

IV. HOLE DOPING EFFECTS

In the section, we discuss a hole doping effect on the system with alternating interactions. At first, we consider the system only with repulsive interactions. It is known that the phase separated magnetically ordered state is realized in the conventional Hubbard model close to half filling at zero temperature.[43] To study how the spatial modulation in the interactions affects the mag-



FIG. 5: (Color online) Density plot of the spontaneous magnetization (a) and the particle density in the sublattice A (b) for the doped system ($n = 0.95$) at $T/D = 0.05$.

netic structure at low temperatures, we calculate the magnetization m_{AF} in the system with $n = 0.95$, as shown in Fig. 5 (a). The magnetization is monotonically decreased by the hole doping, in comparison with the half-filled case (see Fig. 1). It is also found that the phase boundary in the strong coupling region somewhat curves. This behavior should be related to the particle occupation in each sublattice. Figure 5 (b) shows the particle number for the sublattice A. It is found that the commensurability emerges in the sublattice A ($n_A \sim 1$) when $U_A \gg U_B$. In the region, magnetic correlations are enhanced easily, which results in the curved phase boundary. Further hole doping suppresses magnetic correlations, which leads to the phase transition to the paramagnetic state.[43]

By contrast, it is known that the superfluid state is stable against the hole doping. To clarify how the alternating interactions affect the superfluid state, we deal with the attractive Hubbard model with $n = 0.75$ and $U_B/U_A = 5$. Calculating the particle density and pair potential, we obtain the results at fixed temperatures $T/D = 0.02$ and 0.05, as shown in Fig. 6. The paramagnetic metallic state is realized below a certain critical interaction $(U_A)_c$, and the particle number imbalance appears with $n_A \neq n_B$, as shown in Fig. 6 (a). In this case, the Hubbard gap ($\sim |U_B|$) appears in the density of states for the sublattice B, while a low-energy peak clearly appears in the other, as shown in Fig. 7. Beyond the critical interaction $(U_A)_c$, the pair potential Δ is induced and the superfluid state is realized. Then the superfluid gap appears in the vicinity of the Fermi level, as shown in Fig. 7. By examining critical behavior $\Delta \sim |U_A - (U_A)_c|^\beta$ with the exponent $\beta = 1/2$, we determine the critical interactions $(U_A/D)_c = 0.47$ at $T/D = 0.05$ and $(U_A/D)_c = 0.27$ at $T/D = 0.02$, as shown in the inset of Fig. 6 (b). It is also found that the particle number imbalance is suppressed once the system enters the superfluid state, as shown in Fig. 6 (a). Since this tendency is clearly found at lower tempertures, we can say that the realization of the superfluid state makes the system uniform. Similar behavior can be found in the half-filled Hubbard model with a uniform attractive interaction where the hole doping destabilizes the density wave state and the genuine superfluid state is stabilized.



FIG. 6: (Color online) Particle density for each sublattice (a) and pair potential (b) as a function of U_A/D with a fixed ratio $U_B/U_A = 5$. Solid (open) symbols are the results for sublattice $A(B)$ obtained from DMFT with CTQMC method.



FIG. 7: (Color online) Density of states when $U_A/D = -0.4, -0.5$ and -0.75 at $T/D = 0.05$.

When $U_AU_B < 0$, interesting behavior appears in the doped system ($n = 0.75$). The results for the system with a fixed ratio $U_B/U_A = -2$ at $T/D = 0.02$ are shown in Fig. 8. The increase in the interactions monotonically decreases the quasiparticle weight for each sublattice and a singularity appears at $U_A/D \sim 1.7$, which is similar to the half-filled case. However, a further increase in the interaction leads to different behavior. In the sublattice A, the quasiparticle weight becomes almost zero and the local particle density is close to the half filling, as shown in the inset of Fig. 8. Our results suggest the existence of the Mott transition in the sublattice A. In fact, the Mott gap clearly appears in the density of states, as shown in Fig. 9. By contrast, in the sublattice B, the local particle density becomes away from the half filling. Then the quasiparticle weight increases beyond the transition point and the gap structure appears around the Fermi level. This behavior is characteristic of the spin-gap state induced by the pairing transition in the doped system.[44] Therefore, we can say that even in the doped system, the Mott and pairing transitions occur at finite temperatures.



FIG. 8: (Color online) Quasiparticle weight for each sublattice as a function of U_A/D with a fixed ratio $U_B/U_A = -2$. Solid lines with circles are obtained from DMFT with the CTQMC method at $T/D = 0.02$. Dashed lines with squares are obtained from DMFT with the NRG method. Inset shows the particle density for each sublattice.



FIG. 9: (Color online) Density of states for the system with $U_B/U_A = -2$ at $T/D = 0.02$ when $U_A/D = 1.6$ (upper panel) and $1.8D$ (lower panel).

However, the nature of the phase transition at zero temperature may not be trivial since in the system with the uniform interactions, the Mott transition is of second order at half filling and the pairing transition is of first order away from the half filling. To clarify the stability of the normal metallic state, we calculate the renormalization factors at zero temperature by means of the NRG method. We find that the quasiparticle weights smoothly decrease up to $U_A/D \sim 2.0$. In the case, a large difference appears between the NRG and CTQMC results, contrast to the half-filled case. This originates from the fact that the particle number at each sublattice is sensitive to the interaction strength and thermal fluctuations. When the interaction is large enough, oscillation behavior appears in the DMFT iterations and we could not find the stable solution. This may imply that the first-order phase transition occurs and the phase separation is realized at zero temperature, which has been discussed in the attractive Hubbard model.[44] This point thus remains an open issue for future studies.

To clarify the role of the attractive and repulsive interactions in the doped system, we also consider the system with $U_B/U_A = -1/2$, where the magnitude of the attractive interactions is smaller than the other. Fig. 10 shows that when $U_A/D \sim 1$, the renormalization factor in the sublattice A is rapidly decreased, while the other is little decreased. This behavior is contrast to the case $U_B/U_A = -2$ (see Fig. 8), which means the absence of the symmetry axis ($U_A = -U_B$). On the other hand, when the system approaches the phase transition point, the particle number for the sublattice A with repulsive interactions approaches half and both quasiparticle weights decrease. At last, the jump singularity appears in the curves of the quasiparticle weights at $U_A/D \sim 3.3$, where the Mott and pairing transitions occur simultaneously. Therefore, the nature of the phase transition in the doped system is essentially the same as the half-filled case. We wish to note that this is contrast to low-temperature behavior in the doped system with both repulsive interactions. In the system, the stronger repulsive interaction tends to recover the commensurability in the corresponding sublattice, but the weaker one never induces the phase transition in the intermediate band filling. Then the heavy metallic state is realized in the strong coupling regime. Therefore, we can say that a single phase transition discussed here is stabilized by two effects: the Mott transition together with the commensurability due to the repulsive interaction and the pairing transition induced by the attractive interaction.



FIG. 10: (Color online) Quasiparticle weight for each sublattice as a function of U_A/D with a fixed ratio $U_B/U_A = -1/2$. Solid lines with circles are obtained from DMFT with the CTQMC method at $T/D = 0.02$. Dashed lines with squares are obtained from DMFT with the NRG method. Inset shows the particle density for each sublattice.

When the particle number is smaller than the quarter filling ($0 < n < 0.5$), it is expected that the commensurability never emerges and no phase transition occurs. To confirm this, we show the renormalization factors in the systems with $n = 0.4$ and 0.6 at $T/D = 0.02$, as shown in Fig. 11. In the case $n = 0.6$, the singularity appears around $U_A/D \sim 1.6$, which suggests the existence of the single phase transition. On the other hand, when $n = 0.4$, no singularity appears in the curves of the renormalization factors. Therefore, in the doped system with $n \leq 0.5$, the commensurability never emerges and no Mott transition occurs.



FIG. 11: (Color online) Quasiparticle weight for each sublattice as a function of U_A/D with a fixed ratio $U_B/U_A = -2$ at $T/D = 0.02$. Circles (triangles) are the results for $n = 0.6(0.4)$, obtained from DMFT with the CTQMC method.

In this paper, we have discussed how the spatially-modulated interactions affect low-temperature properties in the optical lattice system. We have determined the rich phase diagram of the system. Although this model may be one of the realizable toy models, we have clarified the role of the repulsive and attractive interactions. In particular, the repulsive interaction plays a crucial role in stabilizing the metal-insulator transition in the doped system. This is similar to the essence of the orbital-selective Mott transition in the doped multiband system,[45,46] where the commensurability in the narrower band emerges due to the repulsive interactions. It is an interesting problem whether or not such phase transition indeed occurs in the finite dimensions, which is now under consideration.

V. SUMMARY

We have investigated the fermionic optical lattice system, which is described by the Hubbard model with alternating on-site interactions. By combining dynamical mean-field theory with the continuous-time Monte Carlo method, we have studied low-temperature properties systematically, calculating the quasiparticle weight, double occupancy, and order parameters for each sublattice. It has been clarified that the magnetically ordered state is realized in the system with only repulsive interactions, while the superfluid state is realized in the attractive case. We have also found that when the repulsive and attractive interactions alternate, the metal-insulator transitions occur against magnetic and superfluid fluctuations. In the doped system, we have found that the commensurability emerges due to the repulsive interactions and the insulating state remains with $n > 1/2$.

Acknowledgments

This work was partly supported by the Global Center of Excellence Program "Nanoscience and Quantum Physics" from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan. The simulations have been performed using some of the ALPS libraries.[47]

---

[1] I. Bloch, Nat. Phys. **1**, 23 (2005).
[2] M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch and I. Bloch, Nature **415**, 39 (2002).
[3] R. Jördens, N. Strohmaier, K. Günter, H. Moritz, T. Esslinger, Nature **455**, 204 (2008).
[4] U. Schneider, L. Hackermuller, S. Will, Th. Best, I. Bloch, T. A. Costi, R. W. Helmes, D. Rasch and A. Rosch, Science **322**, 1520 (2008).
[5] S. Jochim,M. Bartenstein, A. Altmeyer, G. Hendl, S. Riedl, C. Chin, J. Hecker Denschlag, and R. Grimm: Science **302** (2003) 2101.
[6] M. W. Zwierlein,C. A. Stan, C. H. Schunck, S. M. F. Raupach, S. Gupta, Z. Hadzibabic, and W. Ketterle: Phys. Rev. Lett. **91** (2003) 250401.
[7] T. Bourdel,L. Khaykovich, J. Cubizolles, J. Zhang, F. Chevy, M. Teichmann, L. Tarruell, S. J. J. M. F. Kokkelmans, and C. Salomon: Phys. Rev. Lett. **93** (2004) 050401.
[8] R. Yamazaki, S. Taie, S. Sugawa, and Y. Takahashi, Phys. Rev. Lett. **105**, 050405 (2010).
[9] T. Saitou, A. Koga, and A. Yamamoto, arXiv:1204.4188.
[10] A. Koga, T. Higashiyama, K. Inaba, S. Suga, and N. Kawakami, J. Phys. Soc. Jap. **77**, 073602 (2008); Phys. Rev. A **79**, 013607 (2009).
[11] M. Snoek, I. Titvinidze, C. Töke, K. Byczuk, and W. Hofstetter, New J. Phys. **10**, 093008 (2008).
[12] A. Georges, G. Kotliar, and W. Krauth, Z. Phys. B **92**, 313 (1993).
[13] M. Caffarel and W. Krauth, Phys. Rev. Lett. **72**, 1545 (1994).
[14] Y. Ohashi and Y. Ono, J. Phys. Soc. Jpn. **70**, 2989 (2001).
[15] W. Metzner and D. Vollhardt, Phys. Rev. Lett. **62**, 324 (1989).
[16] E. Müller-Hartmann, Z. Phys. B: Condens. Matter **74**, 507 (1989).

[17] Th. Pruschke, M. Jarrell, and Freericks, Adv. Phys. **44**, 187 (1995).

[18] A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg, Rev. Mod. Phys. **68**, 13 (1996).

[19] G. Kotliar and D. Vollhardt, Phys. Today **53**, (2004).

[20] E. Gull, A. J. Millis, A. I. Lichtenstein, A. N. Rubtsov, M. Troyer, and P. Werner: Rev. Mod. Phys. **83** (2011) 349.

[21] H. Shiba: Prog. Theor. Phys. **48** (1972) 2171.

[22] A. Yamamoto, S. Yamada, and M. Machida, unpublished.

[23] R. Chitra and G. Kotliar, Phys. Rev. Lett. **83**, 2386 (1999).

[24] A. Georges and G. Kotliar, Phys. Rev. B **45**, 6479 (1992).

[25] X. Y. Zhang, M. J. Rozenberg, and G. Kotliar, Phys. Rev. Lett. **70**, 1666 (1993).

[26] P. Werner, A. Comanac, L. de' Medici, M. Troyer, and A. J. Millis, Phys. Rev. Lett. **97** 076405 (2006).

[27] A. Koga and P. Werner, J. Phys. Soc. Jpn. **79**, 064401 (2010).

[28] N. Takemori and A. Koga, J. Phys. Soc. Jpn. **81**, 063002 (2012).

[29] A. Koga and P. Werner, J. Phys. Soc. Jpn. **79**, 114401 (2010).

[30] D. J. Luitz and F. F. Assaad: Phys. Rev. B **81**, 024509 (2010).

[31] J. Otsuki, H. Kusunose, P. Werner, and Y. Kuramoto: J. Phys. Soc. Jpn. **76** (2007) 114707; J. Otsuki, H. Kusunose, and Y. Kuramoto: Phys. Rev. Lett. **102** (2009) 017202.

[32] F. F. Assaad and T. C. Lang: Phys. Rev. B **76** (2007) 035116; P. Werner and A. J. Millis: Phys. Rev. Lett. **99** (2007) 146404.

[33] A. Koga and P. Werner, Phys. Rev. A **84**, 023638 (2011).

[34] R. Bulla, T. A. Costi, and Th. Pruschke, Rev. Mod. Phys. **80**, 395 (2008).

[35] A. Koga, N. Kawakami, R. Peters, and T. Pruschke, Phys. Rev. B **77**, 045120 (2008); J. Phys. Soc. Jpn. **77**, 033704 (2008).

[36] F. B. Anders and A. Schiller, Phys. Rev. B **74**, 245113 (2006).

[37] R. Peters, Th. Pruschke, and F. B. Anders, Phys. Rev. B **74**, 245114 (2006).

[38] A. Weichselbaum and J. von Delft, Phys. Rev. Lett. **99**, 076402 (2007).

[39] S. F. Gull, in *Maximum Entropy and Bayesian Methods in Science and Engineering*, ed. G. J. Erickson and C. R. Smith (Kluwer Academic, Dordrecht, 1988) p. 53; J. Skilling(Kluwer Academic, Dordrecht, 1989) p. 45; S. F. Gull, *ibid.* p. 53.

[40] R. N. Silver, D. S. Sivia and J. E. Gubernatis, Phys. Rev. B **41**, 2380 (1990); J. E. Gubernatis, M. Jarrell, R. N. Silver and D. S. Sivia, Phys. Rev. B **44**, 6011 (1991).

[41] W. F. Press, S. A. Teukolsky, W. T. Vetterling and B. R. Flannery, *Numerical Recipes* (Cambridge University Press, Cambridge, 1992) p. 809.

[42] R. Bulla and M. Potthoff, Eur. Phys. J. B **13**, 257 (2000).

[43] R. Zitzler, Th. Pruschke, and R. Bulla, Eur. Phys. J. B **27**, 473 (2002).

[44] M. Capone, C. Castellani, and M. Grilli, Phys. Rev. Lett. **88**, 126403 (2002).

[45] A. Koga, N. Kawakami, T. M. Rice, and M. Sigrist, Phys. Rev. Lett. **92**, 216402 (2004).

[46] K. Inaba and A. Koga, J. Phys. Soc. Jpn. **76**, 094712 (2007).

[47] A. F. Albuquerque, F. Alet, P. Corboz, P. Dayal, A. Feiguin, S. Fuchs, L. Gamper, E. Gull, S. Gurtler, A. Honecker, R. Igarashi, M. Korner, A. Kozhevnikov, A. Lauchli, S. R. Manmana, M. Matsumoto, I. P. McCulloch, F. Michel, R. M. Noack, G. Paw.owski, L. Pollet, T. Pruschke, U. Schollwock, S. Todo, S. Trebst, M. Troyer, P. Werner, S. Wessel, J. Magn. Magn. Mater. **310**, 1187 (2007).